

Gastric Bypass Versus Sleeve Gastrectomy

Patient Selection and Short-term Outcome of 47,101 Primary Operations from the Swedish, Norwegian, and Dutch National Quality Registries

Youri Q. M. Poelemeijer, MD,*† Ronald S. L. Liem, MD,‡§ Villy Våge, MD, PhD,|| Tom Mala, MD, PhD,¶
Magnus Sundbom, MD, PhD,# Johan Ottosson, MD, PhD,** and Simon W. Nienhuijs, MD, PhD††

Objective: The aim of this study was to compare the use and short-term outcome of Roux-en-Y gastric bypass (RYGB) and sleeve gastrectomy (SG) in Sweden, Norway, and the Netherlands.

Background: Although bariatric surgery is performed in high volumes worldwide, no consensus exists regarding the choice of bariatric procedure for specific groups of patients.

Methods: Data from 3 national registries for bariatric surgery were used. Patient selection, perioperative data (severe complications, mortality, and rate of readmissions within 30 days), and 1-year results (follow-up rate and weight loss) were studied.

Results: A total of 47,101 primary operations were registered, 33,029 (70.1%) RYGB and 14,072 (29.9%) SG. Patients receiving RYGB met international guidelines for having bariatric surgery more often than those receiving SG (91.9% vs 83.0%, $P < 0.001$). The 2 procedures did not differ in the rate of severe complications (2.6% vs 2.4%, $P = 0.382$), nor 30-day mortality (0.04% vs 0.03%, $P = 0.821$). Readmission rates were higher after RYGB (4.3% vs 3.4%, $P < 0.001$).

One-year post surgery, less RYGB-patients were lost-to follow-up (12.1% vs 16.5%, $P < 0.001$) and RYGB resulted in a higher rate of patients with total weight loss of more than 20% (95.8% vs 84.6%, $P < 0.001$). While the weight-loss after RYGB was similar between hospitals, there was a great variation in weight loss after SG.

Conclusion: This study reflects the pragmatic use and short-term outcome of RYGB and SG in 3 countries in North-Western Europe. Both procedures were safe, with RYGB having higher weight loss and follow-up rates at the cost of a slightly higher 30-day readmission rate.

Keywords: %TWL, bariatric, comparison, complications, DATO, DICA, Dutch, gastrectomy, gastric bypass, LTFU, Netherlands, Norway, Norwegian, outcome, outcomes, quality, readmission, Roux-en-Y, RYGB, SG, sleeve, SOReg, surgery, Sweden, Swedish, total weight loss, TWL

(*Ann Surg* 2019;xx:xxx–xxx)

To ensure and improve the quality of bariatric surgery, assessment of outcome is required.^{1–5} The external validity of case series,

observational studies, and randomized controlled trials may not reflect everyday practice and outcome.⁶ The Rome Diabetes Surgery Summit achieved consensus for a need of standard national registries to collect “real-world” data.^{6,7} In recent years, several national bariatric registries have been established in Europe and early results have been published.^{3,6,8,9}

Although bariatric surgery has been performed in high volumes worldwide for several decades,¹⁰ no consensus exists in regard to the choice of procedure for specific groups of patients.^{11,12} Two procedures dominate at present, Roux-en-Y gastric bypass (RYGB)¹³ and sleeve gastrectomy (SG).^{14,15}

Nationwide registries, suitable for international benchmarking, have been developed in Sweden, Norway, and the Netherlands.^{3,16} A main reason for using data from national registries is to reduce the risk of selection bias and intention-to-treat confounders from individual hospitals. However, selection biases may still impact findings such as choice of surgical procedure for the individual patient, which may to some extent rely on the surgeon’s preference or institutional practice. Moreover, individual countries apply clinical protocols incorporating differences that may impact outcome after bariatric surgery. National differences between individual hospitals may thus be small, while differences in outcome between nations may be more easily found. A recent study showed an extended overview of the registered variables in both registries and also showed that the definitions used for perioperative measures in the registries were comparable thus facilitating evaluation of surgical and outcome indicators between registries.¹⁷ The 30-day morbidity and mortality following primary bariatric surgery in Sweden, Norway, and the Netherlands documented comparable outcome in these countries.¹⁷

The aim of this study was to compare the use and short-term outcome of RYGB and SG in Sweden, Norway, and the Netherlands during 2015 to 2017.

METHODS

Data Registries

Patients receiving a primary RYGB or a primary SG from January 2015 till December 2017 were included. Perioperative data included a 30 days follow-up period after primary surgery. Evaluations of 1-year follow-up outcome included patients operated from January 2015 till December 2016. Patients operated in Sweden, Norway, and the Netherlands were registered in the respective national registries. Patient data were retrieved during the preoperative consultations, during hospital stay, and from follow-up consultations by direct plotting the information into the online-based registries by responsible health care providers. Further details on the design of these registrations have been described previously.^{3,16–19} All procedures open or laparoscopic were included in the registries and analyzed in the present study.

From the *Dutch Institute for Clinical Auditing, Scientific Bureau, Leiden, Netherlands; †Leiden University Medical Center, Department of Surgery, Leiden, Netherlands; ‡Groene Hart Hospital, Department of Surgery, Gouda, Netherlands; §Dutch Obesity Clinic, The Hague, Netherlands; ||Scandinavian Obesity Surgery Registry, Bergen, Norway; ¶Oslo University Hospital, Department of Gastrointestinal Surgery, Oslo, Norway; #Uppsala University, Department of Surgical Sciences, Uppsala, Sweden; **Örebro University Hospital, Department of Surgery, Örebro, Sweden; and ††Catharina Hospital, Department of Surgery, Eindhoven, Netherlands.

The authors declare no conflicts of interest.

Reprints: Youri Q. M. Poelemeijer, MD, Leiden University Medical Center, Department of Surgery, Albinusdreef 2, 2333 ZA Leiden, Netherlands.
E-mail: Y.Q.M.Poелеmeijer@lumc.nl.

Copyright © 2019 Wolters Kluwer Health, Inc. All rights reserved.

ISSN: 0003-4932/16/XXXX-0001

DOI: 10.1097/SLA.0000000000003279

SOReg

SOReg (Scandinavian Obesity Surgery Registry) was initiated in 2007 in Sweden,³ and is supported by the National Board of Health and Welfare, a government agency under the Ministry of Health and Social Affairs. In the last 5 years, more than 98% of all patients who underwent bariatric surgery in Sweden have been registered in SOReg-Sweden (SOReg-S). In 2014, Norway joined SOReg and received status as a nationwide register in June 2015. The coverage rate based on public hospitals was 64% in 2016 and 73% in 2017.

All variables in SOReg-Norway (SOReg-N) and SOReg-S apply the same definitions and the database platform is identical. An identical system for auditing of data to improve quality has been developed. Both the SOReg-N and SOReg-S registries performed external data verification, indicating high quality of data.²⁰

DATO

The nationwide DATO (Dutch Audit for Treatment of Obesity) registry includes all Dutch bariatric patients. Information is collected through an online survey for all patients. The registry officially started on January 1, 2015, and covers over 99.9% of all patients who undergo bariatric surgery in the Netherlands. Nationwide coverage is enforced via the Association of Surgeons of the Netherlands, the umbrella organization of Dutch health insurers, and the Dutch National Health Care Institute.^{16,18} Recent third-party data verification showed an inclusion rate of 100% for all bariatric hospitals in the Netherlands.

The variables have an overlap of more than 90% between the SOReg and DATO registries.¹⁷ For all the variables used in this study, the definitions of DATO correspond to those of SOReg.¹⁷

Main Outcome Measures and Definitions

Six indicators were used to compare the use and outcome of RYGB and SG; patient eligibility for surgery, severe complications, mortality, readmissions, rate of follow-up at 1 year, and 1-year weight loss. Analyses were performed on merged data as well as on national basis.

Eligibility Criteria for Bariatric Surgery

The National Institutes of Health Consensus Development Conference Statement,²¹ recognized as IFSO-guidelines for bariatric surgery, were practiced in all 3 countries.^{7,22–26} To make a distinction between whether or not patients were operated in agreement with these guidelines, a process indicator was established. The patient had to be between 18 and 65 years at the time of surgery, with a body mass index (BMI) of $\geq 40.0 \text{ kg/m}^2$ or a BMI of $35.0\text{--}40.0 \text{ kg/m}^2$ in combination with at least 1 of the 6 major obesity-related diseases: type 2 diabetes mellitus (T2DM), hypertension, dyslipidemia, obstructive sleep apnea syndrome, gastroesophageal reflux disease, and musculoskeletal pain. These diseases were defined by continuous use of medication and with continuous positive airway pressure in sleep apnoea.^{25,27,28} Currently, patients with BMI of $\geq 30.0 \text{ kg/m}^2$ and T2DM may be offered metabolic surgery on an individual basis, but this criterion was not incorporated in the guidelines during the entire study period.²⁸

Perioperative Results

Postoperative complications were categorized according to the Clavien-Dindo Classification of Surgical Complications (CD) and represent complications within 30 days after primary surgery or during the same hospital stay.²⁹ CD-grade IIIb or higher was classified as a severe complication. CD-grade IIIb represents a complication for which a surgical, endoscopic, and/or radiological intervention was performed under general anesthesia. CD-grade

IV is described as life-threatening complications requiring intensive care (IC) admission. CD-grade V reflects the 30-days mortality rate.²⁹ Readmissions within 30-days after surgery were studied.

One-year Results

Patients without a registered 1-year follow-up after primary surgery were considered as lost-to follow-up. Postoperative weight loss is presented by percentage total weight loss (%TWL), which was defined as: $\frac{\text{preoperative weight} - \text{postoperative weight}}{\text{preoperative weight}} \times 100$.^{5,30}

Statistical Analysis

RYGB and SG were analyzed separately. Differences in regard to patient and treatment characteristics were described using frequency tables. Categorical variables were compared using the χ^2 test with Yates' correction. Statistical significance was set at $P < 0.05$. The use of case-mix adjusted outcomes are controversial and not applied.^{31,32}

Results for all 6 process and outcome indicators were presented using funnel plots with 95% control limits that vary in relation to total number of hospital procedures in the above specified study period.^{17,33}

The funnel plots provide information regarding specific process or outcome measures for individual hospitals in relation to the overall average and in relation to results of other anonymized hospitals. The Y-axis shows the percentage of primary bariatric procedures indicated on the X-axis that met the binomial result as the specific indicator indicates. The Y-axis shows the absolute percentage per hospital (dot) and the number of patients operated, as shown on the x-axis.

R version 3.4.3 (Copyright (C) The R Foundation for Statistical Computing Platform) was used for statistical analysis in combination with the 'Companion to Applied Regression'-package (car 2.1–6), and 'A Grammar of Data Manipulation'-package (dplyr 0.7.4).

This study was approved by the regional ethical committee of Stockholm, Sweden (2013/535–31/5) for SOReg-S. The Regional Committee for medical and health research ethics in South East Norway approved the study (reference number: 2018/1631) for SOReg-N. For this study, no ethical approval or informed consent was required under Dutch law for DATO.¹⁶

RESULTS

From January 2015 till December 2017, a total of 47,101 primary bariatric procedures ($>99\%$ laparoscopic cases) were registered. Of these, 33,029 (70.1%) were RYGB and 14,072 (29.9%) were SG procedures. Patient characteristics per country and combined are given in Table 1.

RYGB was the most commonly applied procedure in Sweden (64.0%) and the Netherlands (77.0%), while in Norway, SG was more common (57.0%, $P < 0.001$). Patients who underwent RYGB had more preoperative comorbidities (73.5% vs 64.3%, $P < 0.001$) than patients receiving SG. Patients operated in the Netherlands had more comorbidities than both Sweden and Norway ($P < 0.001$). Moreover, in Norway and Sweden, gastroesophageal reflux disease was about twice as common in RYGB patients, while no difference between the 2 procedures was seen in the Netherlands (Table 1).

Eligibility Criteria for Bariatric Surgery

In total, 89.2% ($n = 42,030$) of patient met the eligibility criteria for bariatric surgery, 91.9% for the RYGB and 83.0% for SG patients, respectively ($P < 0.001$) (Fig. 1). Twenty-three of 59 (39.0%) hospitals [Sweden (SE): 19/28; Norway (NO): 4/13; Netherlands (NL): 0/18] operated significantly more RYGB patients (Fig. 1B), and 18 of 61 (29.5%) hospitals (SE: 13/28; NO: 5/16; NL: 0/17) operated significantly more SG patients (Fig. 1A), not meeting

TABLE 1. Preoperative Patient Characteristics According to Country and Combined Data

	Netherlands				Norway				Sweden				All				P*
	Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	
Number of procedures	21,055	77.0%	6292	23.0%	1365	43.0%	1809	57.0%	10,609	64.0%	5971	36.0%	33,029	70.1%	14,072	29.9%	<0.001
Patient characteristics																	
Age (mean, y, SD)	44.5	± 10.9	41.9	± 12.4	42.9	± 10.8	42.1	± 11.1	40.7	± 11.8	41.0	± 10.9	43.2	± 11.2	41.5	± 11.6	<0.001
BMI (mean, kg/m ² , SD)	43.5	± 5.0	45.6	± 6.6	43.3	± 5.2	41.9	± 5.4	41.7	± 5.6	39.5	± 5.4	42.9	± 5.2	42.5	± 5.9	<0.001
Female	17,064	81.0%	4684	74.4%	1045	76.6%	1395	77.1%	2188	77.8%	4834	80.1%	26,171	79.2%	10,913	77.6%	<0.001
Comorbidities	17,139	81.4%	5017	79.7%	890	65.2%	1109	61.3%	6256	59.0%	2917	48.9%	24,285	73.5%	9043	64.3%	<0.001
T2DM	4521	21.5%	1058	16.8%	199	14.6%	196	10.8%	1456	13.7%	502	8.4%	6176	18.7%	1756	12.5%	<0.001
Hypertension	7312	34.7%	2016	32.0%	453	33.2%	475	26.3%	2791	26.3%	1,208	20.2%	10,556	32.0%	3699	26.3%	<0.001
Dyslipidemia	4260	20.2%	1060	16.8%	228	16.7%	178	9.8%	1046	9.9%	408	6.8%	5534	16.8%	1646	11.7%	<0.001
GERD	2705	12.8%	784	12.5%	293	21.5%	175	9.7%	1322	12.5%	341	5.7%	4320	13.1%	1300	9.2%	<0.001
OSAS	3841	18.2%	1205	19.2%	223	16.3%	226	12.5%	1113	10.5%	412	6.9%	5177	15.7%	1843	13.1%	<0.001
Musculoskeletal pain	9519	45.2%	2843	45.2%	462	33.8%	545	30.1%	2337	22.0%	877	14.7%	12,318	37.3%	4265	30.3%	<0.001

BMI indicates body mass index; GERD, gastroesophageal reflux disease; N/A, not available; OSAS, obstructive sleep apnea syndrome; T2DM, type 2 diabetes mellitus. *P values compared all 3 countries together.

the eligibility criteria for bariatric surgery compared with the overall average.

Complicated Postoperative Course

Severe complications (CD-Grade ≥IIIb) were registered in 2.6% (n = 846) patients after RYGB and 2.4% (n = 341) patients

after SG (P = 0.382). Reinterventions due to severe complications were performed in 2.0% (n = 667) patients after RYGB and 2.1% (n = 290) patients after SG (P = 0.771). The overall number of patients registered with a CD-grade IV complication was 0.5% (n = 170) for RYGB and 0.3% (n = 47) after SG (P = 0.008) (Table 2). In the RYGB group, 4 of 59 (6.8%) hospitals (SE: 1/28; NO: 0/13; NL: 3/18)




FIGURE 1. Percentage of primary bariatric procedures meeting the inclusion criteria, reported per hospital. C.I. indicates control limit; RYGB, Roux-en-Y gastric bypass procedure; SG, sleeve gastrectomy.

TABLE 2. Morbidity and Mortality After Primary Sleeve Gastrectomy and Gastric Bypass Procedure

	Netherlands				Norway				Sweden				All				P*
	Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		Roux-en-Y Gastric Bypass		Sleeve Gastrectomy		
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	
Number of procedures	21,055	77.0%	6292	23.0%	1365	43.0%	1809	57.0%	10,609	64.0%	5971	36.0%	33,029	70.1%	14,072	29.9%	< 0.001
Perioperative complications	420	2.0%	131	2.1%	66	4.8%	32	1.8%	284	2.7%	57	1.0%	770	2.3%	220	1.6%	< 0.001
Gastrointestinal perforation	135	0.6%	7	0.1%	24	1.8%	1	0.1%	121	1.1%	3	0.1%	280	0.8%	11	0.1%	
Bleeding	64	0.3%	44	0.7%	5	0.4%	0	0.0%	10	0.1%	12	0.2%	79	0.2%	56	0.4%	
Spleen injury	31	0.1%	21	0.3%	4	0.3%	8	0.4%	11	0.1%	17	0.3%	46	0.1%	46	0.3%	
Hepatic injury	44	0.2%	11	0.2%	0	0.0%	0	0.0%	16	0.2%	8	0.1%	60	0.2%	19	0.1%	
Major vascular injury	1	0.0%	1	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	1	0.0%	1	0.0%	
Postoperative complications	1046	5.0%	347	5.5%	80	5.9%	94	5.2%	753	7.1%	309	5.2%	1879	5.7%	750	5.3%	< 0.001
Bleeding	314	1.5%	129	2.1%	25	1.8%	25	1.4%	123	1.2%	66	1.1%	462	1.4%	220	1.6%	
Leakage	109	0.5%	46	0.7%	9	0.7%	16	0.9%	80	0.8%	30	0.5%	198	0.6%	92	0.7%	
Intra-abdominal infection	29	0.1%	15	0.2%	5	0.4%	11	0.6%	64	0.6%	22	0.4%	98	0.3%	48	0.3%	
Wound infection	28	0.1%	11	0.2%	4	0.3%	11	0.6%	54	0.5%	45	0.8%	86	0.3%	67	0.5%	
Intestinal obstruction	63	0.3%	10	0.2%	16	1.2%	0	0.0%	113	1.1%	7	0.1%	192	0.6%	17	0.1%	
Cardiac events	38	0.2%	12	0.2%	0	0.0%	5	0.3%	6	0.1%	3	0.1%	44	0.1%	20	0.1%	
Pulmonary events	71	0.3%	22	0.3%	5	0.4%	1	0.1%	33	0.3%	17	0.3%	109	0.3%	40	0.3%	
Thrombotic events	8	0.0%	4	0.1%	1	0.1%	2	0.1%	9	0.1%	2	0.0%	18	0.1%	8	0.1%	
Other	427	2.0%	115	1.8%	40	2.9%	46	2.5%	327	3.1%	164	2.7%	794	2.4%	325	2.3%	
Overall	506	2.4%	205	3.3%	29	2.1%	31	1.7%	315	3.0%	105	1.8%	850	2.6%	341	2.4%	< 0.001
Reintervention CD-grade IIIb	346	1.6%	163	2.6%	27	2.0%	28	1.5%	294	2.8%	99	1.7%	667	2.0%	290	2.1%	
IC/ICU admission CD-grade IV	148	0.7%	38	0.6%	2	0.1%	3	0.2%	20	0.2%	6	0.1%	170	0.5%	47	0.3%	
Mortality CD-grade V	12	0.1%	4	0.1%	0	0.0%	0	0.0%	1	0.0%	0	0.0%	13	0.0%	4	0.0%	
Length of stay and readmission																	
Readmissions (<30 d)	572	2.7%	158	2.5%	87	4.6%	87	4.8%	752	7.1%	240	4.0%	1411	4.3%	485	3.4%	< 0.001
Hospital stay (mean, d, SD)	1.6 ± 2.9		1.6 ± 2.7		1.7 ± 3.4		2.0 ± 2.2		1.5 ± 4.6		1.6 ± 2.3		1.6 ± 3.5		1.7 ± 2.5		< 0.001

CD indicates Clavien-Dindo Classification; IC, intensive care; ICU, intensive care unit; N/A, not available.

*P values compared all 3 different countries together.

registered significantly lower rate of complications than the average. In contrast, a higher rate of severe complications was seen for SG in 9 of 61 (14.8%) hospitals (SE: 6/28; NO: 1/16; NL: 2/17) (Fig. 2). Thirty-day mortality was 0.04% (n = 13) after RYGB and 0.03% (n = 4) after SG (P = 0.821).

The 3 most common complications after RYGB and SG combined were bleeding (1.6%), leakage (0.7%), and wound infection (0.5%), with no statistical difference between the 2 procedures (Table 2).

Length of Hospital Stay and Readmissions

The length of stay was shorter after RYGB than SG, 1.6 days (SD ± 3.5) and 1.7 days (SD ± 2.5), respectively (P < 0.001) (Table 2). In the Netherlands, the length of hospital stay after RYGB and SG was comparable, while in Norway and Sweden, hospital stay was somewhat shorter after RYGB than SG in the same country (P < 0.001).

The readmission rate was 4.3% (n = 1411) after RYGB and 3.4% (n = 485) after SG (P < 0.001). Readmission rates were lowest in the Netherlands. Significantly more Swedish hospitals registered a readmission after RYGB (7.1%) than the overall average (P < 0.001) (Table 2).

Lost to Follow-up After 1 Year

On average, the 1-year lost to follow-up was lower after RYGB than SG, 12.1% (n = 2712) and 16.5% (n = 1433), respectively (P < 0.001). The largest difference between the 2 procedures was seen in Sweden, 11.9% for RYGB versus 20.1% for SG, respectively (P < 0.001). (Fig. 3).

After RYGB, 12 of 59 (20.3%) (SE: 5/28; NO: 2/13; NL: 5/18) and after SG 8 of 61 (13.1%) (SE: 5/28, NO: 1/16; NL: 2/17) centers

had significantly higher rates of lost to follow-up up than the overall average, respectively.

Total Weight Loss (%TWL)

ATWL percent (%TWL) of more than 20% the first year after surgery was reached more often after RYGB than SG, 95.8% and 84.6% of the patients, respectively (P < 0.001) (Fig. 4). In total, after RYGB, 6 of 59 (10.2%) (SE: 4/28; NO: 1/13; NL: 1/18) hospitals and after SG 18 of 61 (29.5%) (SE: 16/28; NO: 0/16; NL: 2/17) hospitals scored significantly lower than the overall average.

There is a significant difference in 20%TWL after SG in Sweden (75.2%), Norway (93.4%), and the Netherlands (90.8%) (P < 0.001), while the difference is much smaller after RYGB (SE: 94.9%; NO: 95.0%; NL: 96.3%; P < 0.001). The considerable spread in outcomes between hospitals after SG compared with RYGB is easily noticeable in the funnel plot.

DISCUSSION

This study describes the pragmatic everyday outcome of RYGB and SG in 3 Northwest-European countries based on a standard platform of 6 quality indicators. RYGB was more often used in adherence to commonly accepted guidelines for eligibility to bariatric surgery. Postoperative severe complications (2.6% vs 2.4%) and mortality rates (0.04% vs 0.03%) did not differ between RYGB and SG, but readmission rates were higher after RYGB (4.3% vs 3.4%). In RYGB-patients, however, 1-year results were superior, with lower lost-to follow-up (12.1% vs 16.5%) and higher rate of %TWL ≥ 20% (95.8% vs 84.6%). These benchmarking findings may act as guidelines for expected early outcome of bariatric surgery in Northwest-Europe and elsewhere.




FIGURE 2. Percentage of primary bariatric procedures with a severe complication within 30 days after primary surgery, reported per hospital. C.I. indicates control limit; RYGB, Roux-en-Y gastric bypass procedure; SG, sleeve gastrectomy.

Most patients were operated in adherence to internationally used IFSO-guidelines for eligibility to bariatric surgery. Indications for bariatric surgery differ among European countries despite agreement on the international clinical guidelines.³⁴ At the moment, the BMI inclusion criteria in the Netherlands, Sweden, and Norway have been set at a BMI of 40.0 kg/m² or a BMI of 35.0 kg/m² for patients with obesity-related disease. Interestingly, we found that more RYGB patients were operated according to international guidelines for bariatric surgery than SG patients. This could be due the fact that there are some Swedish private clinics performing SG on patients with a BMI of 30 to 35 kg/m² or 35 to 40 kg/m² without any obesity-related comorbidity.

Comparing the 3 countries, the annual hospital case load was highest in the Netherlands. In all countries, mostly females were operated at a rate of about 75% or higher. Dutch patients were significantly older, had a higher BMI, and a higher number of preoperative comorbidities. The mean age and BMI in the present study is lower than reported in most American series.³⁵ Overall, as well as in Sweden and the Netherlands, RYGB was the most common primary procedure, whereas in Norway, SG has reached 57% based on registered data. In the United States, SG surpassed RYGB in 2013³⁶ and was estimated by ASMBS to constitute of 53% to 59% of all bariatric procedures during the present study period (2015 to 2017).³⁷ Interestingly, the findings indicate that there may be between-country differences in regarding to the use of SG in patients with gastroesophageal reflux disease (Table 1).

The present overall rates of severe postoperative complications were in line with results presented in the international literature.

In a recent review on 107,874 patients,³⁵ the leak rate was 1.1% in RYGB and 1.2% in SG. The associated factors for severe postoperative complications are laparoscopic versus open surgery, older age, surgical procedural experience, preoperative comorbidities, and BMI.^{38–41} In Sweden, more postoperative complications were registered for RYGB surgery than SG (7.1% vs 5.2%). As significant outliers from all 3 countries were visible in all measured process and outcome indicators, this may provide insight to areas in need of improvements.

The present mortality of 0.03% to 0.04% compares favorably to others; Gribsholt et al⁴² presenting 0.04% in 9895 Danish RYGBs and the 0.20% to 0.22% presented in a large American cohort (n = 43,354) of RYGB and SG⁴³ and a recent meta-analysis on 69,494 patients.⁴⁴ These results underline the safety of RYGB and SG in these patients.

The overall hospital stay was short for both procedures (1.6 and 1.7 days, respectively), while Norway and Sweden had a significantly higher percentage of readmissions, especially after RYGB (4.6% and 7.1%, respectively). This could partly be explained by demographic differences between countries, where people often live closer to a (bariatric) hospital in the Netherlands. However, it could also be explained by centralization in the Netherlands, where hospitals perform a higher volume of bariatric procedures per year per clinic as demonstrated in Figs. 1 to 4, and therefore have a higher number of procedures per surgeon per year. This could also explain the lower postoperative complication ratio described earlier.

Although it is generally not recommended to report weight loss with less than 2-year follow up, the present 1-year weight loss




FIGURE 3. Percentage of procedures with a registered 12-month follow-up after primary surgery, reported per hospital. C.I. indicates control limit; RYGB, Roux-en-Y gastric bypass; SG, sleeve gastrectomy.

after RYGB is similar and more predictable between institutions, although there is a great variation in weight loss after SG. This can reflect the need for high technical quality in both procedures, as patients receiving large pouches and sleeves are known to have inferior weight loss. According to our data, it thus seems more difficult to achieve a technical perfect SG. The presented weight loss is in line with the literature as well as earlier results from SOReg-S on patients operated before 2014 in Sweden.^{17,19} Interestingly, the 1-year %TWL outcome in RYGB seems more uniform and predictable across hospitals and nations than that of SG.

Currently, almost 98% of elective bariatric surgical cases are managed worldwide by laparoscopy.¹² The increased use of laparoscopy in bariatric surgery has reduced postoperative morbidity and mortality.⁴⁵⁻⁴⁷ In an attempt to further improve quality, a minimum annual procedural volume per hospital of 200 cases has been established in the Netherlands and 100 in Sweden. In Norway, no formal minimum annual procedural hospital volume exists, although most centers perform close to 100 or more procedures annually. Although numerous studies suggest an inverse relation between hospital case load volume and postoperative severe complicated course, recent studies show no significant benefits for choosing a high-volume hospital compared with a low-volume hospital.^{38-40,48,49} This remains a topic of discussion and accreditation on quality outcomes may be of greater value than that of volume.

Several aspects of long-term outcomes can be monitored in Sweden and Norway by registering the unique identification number given at birth, which allows cross-linking with other nationwide registrations. All Dutch citizens have a similar identification number,

but cross-linking between different registries may be more challenging due to legal restrictions. However, the national bariatric registries can learn from one another and allow for international comparison. The SOReg registries include more pre- and postoperative laboratory values, which allow comorbidities to be measured more objectively. On the contrary, registration in DATO is mandatory, whereas registration in SOReg-N is on voluntary basis. In addition, the Netherlands offers a weekly benchmark feedback, while SOReg presents a selection of quality outcomes online and publish annual figures.

Case-mix adjusted outcomes are still controversial in the international literature^{31,32} and deliberately not applied in this study and could imply a possible selection bias as does the heterogeneity of the data, which reflects real-world data. This could be a limitation of this study in a narrower sense of the word.

Nationwide clinical audits provide detailed information on patient characteristics, treatment, and individual hospitals. This information is quickly available for monitoring of quality indicators. These indicators can be used for hospitals to compare their performances relative to a national and an international benchmark analysis. A disadvantage of national clinical audits is that data may not always be complete and directly validated.

The major strength of this study is the international, population-based design, the use of pooled data from 3 high-quality registries including in-depth information, and almost complete coverage of all patients who had bariatric surgery in the Netherlands and Sweden. Standardization of registries and consensus of definitions of measures included facilitate comparisons between countries that may impact quality of the treatment given on an international level.




FIGURE 4. Percentage of primary bariatric procedures registered with %TWL ≥ 20 12 months after primary surgery, reported per hospital. C.I. indicates control limit; RYGB, Roux-en-Y gastric bypass procedure; SG, sleeve gastrectomy.

ACKNOWLEDGMENT

The authors thank all surgeons, registrars, physician assistants, and administrative nurses who registered patients in the DATO, SOReg-N, and SOReg-S.

REFERENCES

1. Van Leersum NJ, Snijders HS, Henneman D, et al. The Dutch surgical colorectal audit. *Eur J Surg Oncol.* 2013;39:1063–1070.
2. Hopkins JC, Howes N, Chalmers K, et al. Outcome reporting in bariatric surgery: an in-depth analysis to inform the development of a core outcome set, the BARIACT Study. *Obes Rev.* 2015;16:88–106.
3. Hedenbro JL, Naslund E, Boman L, et al. Formation of the Scandinavian Obesity Surgery Registry, SOReg. *Obes Surg.* 2015;25:1893–1900.
4. Coulman KD, Abdelrahman T, Owen-Smith A, et al. Patient-reported outcomes in bariatric surgery: a systematic review of standards of reporting. *Obes Rev.* 2013;14:707–720.
5. Brethauer SA, Kim J, el Chaar M, et al. Standardized outcomes reporting in metabolic and bariatric surgery. *Surg Obes Relat Dis.* 2015;11:489–506.
6. Hopkins J, Welbourn R. The importance of national registries/databases in metabolic surgery: the UK experience. *Surg Obes Relat Dis.* 2016;12:1178–1185.
7. Rubino F, Kaplan LM, Schauer PR, et al., Diabetes Surgery Summit D. The Diabetes Surgery Summit consensus conference: recommendations for the evaluation and use of gastrointestinal surgery to treat type 2 diabetes mellitus. *Ann Surg.* 2010;251:399–405.
8. Stroh C, Weiner R, Horbach T, et al. [New data on quality assurance in bariatric surgery in Germany]. *Zentralbl Chir.* 2013;138:180–188.
9. Debs T, Petrucciani N, Kassir R, et al. Trends of bariatric surgery in France during the last 10 years: analysis of 267,466 procedures from 2005–2014. *Surg Obes Relat Dis.* 2016;12:1602–1609.
10. Angrisani L, Santonicola A, Iovino P, et al. Bariatric surgery worldwide 2013. *Obes Surg.* 2015;25:1822–1832.
11. Angrisani L, Santonicola A, Iovino P, et al. Bariatric surgery and endoluminal procedures: IFSO worldwide survey 2014. *Obes Surg.* 2017;27:2279–2289.
12. Welbourn R, Pournaras DJ, Dixon J, et al. Bariatric surgery worldwide: baseline demographic description and one-year outcomes from the Second IFSO Global Registry Report 2013–2015. *Obes Surg.* 2018;28:313–322.
13. Mason EE, Ito C. Gastric bypass. *Ann Surg.* 1969;170:329–339.
14. Lee WJ, Almulaifi AM, Tsou JJ, et al. Duodenal-jejunal bypass with sleeve gastrectomy versus the sleeve gastrectomy procedure alone: the role of duodenal exclusion. *Surg Obes Relat Dis.* 2015;11:765–770.
15. Lager CJ, Esfandiari NH, Subauste AR, et al. Roux-En-Y gastric bypass vs. sleeve gastrectomy: balancing the risks of surgery with the benefits of weight loss. *Obes Surg.* 2017;27:154–161.
16. Poelmeijer YQM, Liem RSL, Nienhuijs SW. A Dutch Nationwide Bariatric Quality Registry: DATO. *Obes Surg.* 2018;28:1602–1610.
17. Poelmeijer YQM, Liem RSL, Vage V, et al. Perioperative outcomes of primary bariatric surgery in North-Western Europe: a pooled multinational registry analysis. *Obes Surg.* 2018;28:3916–3922.
18. Jaarrapport 2016. Dutch Institute for Clinical Auditing; 2017 [08–2017]. Available at: <https://dica.nl/jaarrapportage-2016>.
19. Årsrapport 2016. Scandinavian Obesity Surgery Registry (SOReg); 2017 [cited January 10, 2017]. Available at: <http://www.ucr.uu.se/soreg/>.
20. Tao W, Holmberg D, Naslund E, et al. Validation of obesity surgery data in the Swedish National Patient Registry and Scandinavian Obesity Registry (SOReg). *Obes Surg.* 2016;26:1750–1756.
21. Gastrointestinal surgery for severe obesity: National Institutes of Health Consensus Development Conference Statement. *Am J Clin Nutr* 1992; 55(2 suppl):615S–619S.
22. Bays HE, Jones PH, Jacobson TA, et al. Lipids and bariatric procedures part 1 of 2: scientific statement from the National Lipid Association, American

- Society for Metabolic and Bariatric Surgery, and Obesity Medicine Association: full report. *J Clin Lipidol*. 2016;10:33–57.
23. Mechanick JI, Kushner RF, Sugerman HJ, et al. American Association of Clinical Endocrinologists, The Obesity Society, and American Society for Metabolic & Bariatric Surgery medical guidelines for clinical practice for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient. *Obesity (Silver Spring)*. 2009;17(suppl 1):S1–S70.
 24. Poirier P, Cornier MA, Mazzone T, et al. Bariatric surgery and cardiovascular risk factors: a scientific statement from the American Heart Association. *Circulation*. 2011;123:1683–1701.
 25. Sauerland S, Angrisani L, Belachew M, et al. Obesity surgery: evidence-based guidelines of the European Association for Endoscopic Surgery (EAES). *Surg Endosc*. 2005;19:200–221.
 26. Melissas J. IFSO guidelines for safety, quality, and excellence in bariatric surgery. *Obes Surg*. 2008;18:497–500.
 27. Paulus GF, de Vaan LE, Verdam FJ, et al. Bariatric surgery in morbidly obese adolescents: a systematic review and meta-analysis. *Obes Surg*. 2015;25:860–878.
 28. Fried M, Yumuk V, Oppert JM, et al. Interdisciplinary European guidelines on metabolic and bariatric surgery. *Obes Surg*. 2014;24:42–55.
 29. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240:205–213.
 30. van de Laar A, de Caluwe L, Dillemans B. Relative outcome measures for bariatric surgery. Evidence against excess weight loss and excess body mass index loss from a series of laparoscopic Roux-en-Y gastric bypass patients. *Obes Surg*. 2011;21:763–767.
 31. DeMaria EJ, Murr M, Byrne TK, et al. Validation of the obesity surgery mortality risk score in a multicenter study proves it stratifies mortality risk in patients undergoing gastric bypass for morbid obesity. *Ann Surg*. 2007;246:578–582. discussion 83–84.
 32. Garcia-Garcia ML, Martin-Lorenzo JG, Liron-Ruiz R, et al. Failure of the Obesity Surgery Mortality Risk Score (OS-MRS) to predict postoperative complications after bariatric surgery. A single-center series and systematic review. *Obes Surg*. 2017;27:1423–1429.
 33. Spiegelhalter DJ. Funnel plots for comparing institutional performance. *Stat Med*. 2005;24:1185–1202.
 34. Borisenko O, Colpan Z, Dillemans B, et al. Clinical indications, utilization, and funding of bariatric surgery in Europe. *Obes Surg*. 2015;25:1408–1416.
 35. Chang SH, Freeman NLB, Lee JA, et al. Early major complications after bariatric surgery in the USA, 2003–2014: a systematic review and meta-analysis. *Obes Rev*. 2018;19:529–537.
 36. Spaniolas K, Kasten KR, Brinkley J, et al. The changing bariatric surgery landscape in the USA. *Obes Surg*. 2015;25:1544–1546.
 37. American Society for Metabolic and Bariatric Surgery. Estimate of Bariatric Surgery Numbers, 2011–2017. American Society for Metabolic and Bariatric Surgery; 2018. Available at: <https://asmbs.org/resources/estimate-of-bariatric-surgery-numbers>. Accessed May 11, 2018.
 38. Ricci C, Gaeta M, Rausa E, et al. Long-term effects of bariatric surgery on type II diabetes, hypertension and hyperlipidemia: a meta-analysis and meta-regression study with 5-year follow-up. *Obes Surg*. 2015;25:397–405.
 39. Rosenthal RJ, Szomstein S, Kennedy CI, et al. Laparoscopic surgery for morbid obesity: 1,001 consecutive bariatric operations performed at The Bariatric Institute, Cleveland Clinic Florida. *Obes Surg*. 2006;16:119–124.
 40. Parikh MS, Laker S, Weiner M, et al. Objective comparison of complications resulting from laparoscopic bariatric procedures. *J Am Coll Surg*. 2006;202:252–261.
 41. Stenberg E, Szabo E, Agren G, et al. Early complications after laparoscopic gastric bypass surgery: results from the Scandinavian Obesity Surgery Registry. *Ann Surg*. 2014;260:1040–1047.
 42. Gribsholt SB, Thomsen RW, Svensson E, et al. Overall and cause-specific mortality after Roux-en-Y gastric bypass surgery: a nationwide cohort study. *Surg Obes Relat Dis*. 2017;13:581–587.
 43. Chaar ME, Lundberg P, Stoltzfus J. Thirty-day outcomes of sleeve gastrectomy versus Roux-en-Y gastric bypass: first report based on Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program database. *Surg Obes Relat Dis*. 2018;14:545–551.
 44. Rausa E, Bonavina L, Asti E, et al. Rate of death and complications in laparoscopic and open Roux-en-Y gastric bypass. A meta-analysis and meta-regression analysis on 69,494 patients. *Obes Surg*. 2016;26:1956–1963.
 45. Lujan JA, Frutos MD, Hernandez Q, et al. Laparoscopic versus open gastric bypass in the treatment of morbid obesity: a randomized prospective study. *Ann Surg*. 2004;239:433–437.
 46. Siddiqui A, Livingston E, Huerta S. A comparison of open and laparoscopic Roux-en-Y gastric bypass surgery for morbid and super obesity: a decision-analysis model. *Am J Surg*. 2006;192:e1–e7.
 47. Tian HL, Tian JH, Yang KH, et al. The effects of laparoscopic vs. open gastric bypass for morbid obesity: a systematic review and meta-analysis of randomized controlled trials. *Obes Rev*. 2011;12:254–260.
 48. Jafari MD, Jafari F, Young MT, et al. Volume and outcome relationship in bariatric surgery in the laparoscopic era. *Surg Endosc*. 2013;27:4539–4546.
 49. Jonker FHW, Hagemans JAW, Burger JWA, et al. The influence of hospital volume on long-term oncological outcome after rectal cancer surgery. *Int J Colorectal Dis*. 2017;32:1741–1747.